

EARLY LEVEL 1B VALIDATION READINESS

PRESENTATION TO AIRS SCIENCE TEAM

FEBRUARY 12, 2002

J. SUSSKIND, C. BARNET, J. BLAISDELL, L. IREDELL,
L. KOUVARIS, F. KEITA

Early Level 1B Validation Readiness

Things We Proposed To Do

- Examine earliest AIRS spectra for reasonableness

- Compute AIRS radiance biases and standard deviations for clear cases

- Generate channel noise covariance matrix (with and without tuning)

- Produce HIRS3 like radiances from AIRS and generate TOVS Pathfinder like retrievals

Are We Ready?

- All areas need work

- New computing hardware needed

 - Readiness contingent on timely delivery

Examine Earliest AIRS Spectra For Reasonableness

Compute AIRS Biases And Standard Deviations For Clear Cases

Method

- Compute spectral differences between observed radiances and computed radiances for clear cases
- Use NCEP forecast to compute radiances for 1 whole day of observations

Need to Install and Integrate

- Larry's angle correction
- Mitch's clear test
- Ability to read and colocate NCEP forecast (from Mitch)
- Code to fill in everything needed to compute radiances (from Mitch)

Need to Develop

- Code to compute statistics of radiance differences as a function of zenith angle
 - Mean and standard deviations
- Code to display zenith angle dependence of residuals for select channels

None of these are very difficult and we should be ready

There are 324000 retrievals per day

- If 3% called "clear" get 9720 clear cases per day
- Roughly 300 cases per zenith angle for 1 day

Should be ready by launch + 3 months

Generate Channel Noise Covariance Matrix

$\Sigma_{\text{NCHAN,NCASE}} = R_{\text{OBS}} - R_{\text{COMP}} =$ Matrix of residuals

$\frac{1}{\text{NCHAN}} (\Sigma_{\text{OBS}} - \Sigma_{\text{COMP}}) (\Sigma_{\text{OBS}} - \Sigma_{\text{COMP}})^T$ $\Sigma_{\text{NCHAN,NCHAN}} =$ Noise covariance matrix

If we remove bias only, channel noise covariance remains unchanged

“Tuning” may remove bias and lower noise covariance

$$R_{\text{COMP}} = R_{\text{COMP}} + \Sigma_{\text{COMP}}$$

We will investigate simple forms of Σ_{COMP}

$$\Sigma_{\text{COMP}} = A + B(M_{\text{OBS}} - \bar{M}_{\text{OBS}}) + C(1 - \cos \theta) + D \sin(\theta)$$

M_{OBS} is matrix of AMSU and HSB channel observations

See if covariance of (Σ_{OBS}) is lower

See which channel errors are correlated

None of the above is in place

Should be ready by launch + 3 months

Produce HIRS3 Like Radiances From AIRS And Generate TOVS Pathfinder-Like Retrievals

Method

Simulate HIRS3 radiances from AIRS simulated radiances

$$R_{I,HIRS} = \sum A_{IJ} R_{J,AIRS}$$

Use granule 401 simulation to generate coefficients

Simulate HIRS3 data from AIRS observed data

Modify Pathfinder program to incorporate simulated HIRS and AMSU data

Run TOVS Pathfinder program for 1 month with data simulated from AIRS observations

Compare results with NOAA 16 retrievals

Should be similar if AIRS is performing well

Needs a lot of preparation

May be ready by launch + 3 months

